

OSTEOLOGICAL EVALUATION

Prepared by
EVAN MATSHES BSc, MD
Consultant Osteologist



Product No. BC-209

**Human 16-month-old
Child Skull
(16-18 months)**



Bone Clones, Inc.

OSTEOLOGICAL REPRODUCTIONS

9200 Eton Ave. Chatsworth, CA 91311

Phone: (818) 709-7991 or (800) 914-0091 (USA only)

Email: info@boneclones.com Web: www.boneclones.com

© Bone Clones, Inc. 2015

Human, Child (16-18 months)

Product Number: BC-209

Specimen Evaluated: Bone Clones® replica

Skeletal Inventory: 1 intact cranium
1 intact mandible

General observations:

In general, the molding process has preserved significant details necessary for evaluation. The general shape and configuration of the skull is within normal limits. The general morphology of the individual visible cranial bones is within normal limits. Sutural patterns are of expected configuration. There is the (nasal) remnant of a partial metopic suture. Remnants of the mendosal suture are at the right and left lateral extents of the occipital bone. The anterior and posterior intra-occipital sutures are not fused. There is a right epipteric bone and the suggestion of a left epipteric bone. The foramina are of expected configuration. The skull is atraumatic.

The floors of the external auditory meati are open bilaterally. Multiple ectocranial vascular grooves course over the surface of the skull. An irregular, 1.4 cm diameter pitted depression is in the posterior paramidline occipital bone at the level of the mendosal suture. There is an irregular 1.1 x 0.7 cm patterned depression of the left parietal bone immediately overlying the midline aspect of the squamous suture.

Dentition:

There are 8 teeth in the maxillary arcade and 9 teeth in the mandibular arcade. All teeth have a deciduous morphology; there are no adult teeth.

The following fully erupted teeth are present in the maxillae: 5.4 [B], 5.2 [D], 5.1 [E], 6.1 [F], 6.2 [G], and 6.4 [I].

The following fully erupted teeth are present in the mandible: 7.4 [L], 7.2 [N], 7.1 [O], 8.1 [P], and 8.4 [S].

The following partially erupted teeth are present in the maxillae: 5.3 [C] and 6.3 [H].

The following partially erupted teeth are present in the mandible: 7.3 [M] and 8.3 [R].

Bone Clones ® Descriptive Analysis Report

The following teeth are beginning to erupt through the maxillary cortical bone: 5.5 [A] and 6.5 [J].

The following teeth are beginning to erupt through the mandibular cortical bone: 7.5 [K] and 8.5 [T].

The following unerupted (but exposed) teeth have evidence of calcification: 1.6 [#3], 2.6 [#14], 3.6 [#19], and 4.6 [#30].

The atraumatic gomphosis of 8.2 [Q] is empty and is without signs of healing.

There is an irregular, bulbous projection on the left anterior palate.

There are no dental restorations or prostheses. There is no significant attrition.

Non-Dental Features of Age:

Fontanelles

The anterior fontanelle is open; it is 1.1 cm in the anteroposterior plane, and 1.4 cm in the transverse plane. The posterior, sphenoidal (anterolateral) and mastoidal (posterolateral) fontanelles are closed.

The speno-occipital synchondrosis is open.

The calvarial sutures are all open (there is no evidence of ossification).

SUMMARY:

1. Age

Dental

Likely 16 – 18 months.

Non-Dental

Likely less than 26 months, greater than 12 months.

Anterior fontanelle open.

Closure: median 13.8 months[1], range 4 – 26 months.[2]

Posterior fontanelle closed.

Closure: 2 – 3 months.[3]

Sphenoidal (anterolateral) fontanelle closed.

Closure: 2-3 months.[3]

Mastoidal (posterolateral) fontanelle closed.

Closure: 1 year.[3]

Spheno-occipital synchondrosis open.

Closure: 10.5 – 16 years.[4, 5]

Posterior intra-occipital suture open.

Closure: 1 – 3 years.[6]

Anterior intra-occipital suture open.

Closure: 5 – 7 years.[6]

EDUCATIONAL RESOURCES:

1. This is an excellent example of a young child's skull.
2. It may be appropriate to discuss the differences between primary and secondary dentition, eruption patterns, and controversies surrounding the timelines that 'typify' those eruption patterns.
3. Age assessment of skeletal remains is best done in the context of the entire skeleton. It is important for educators to emphasize that when limited to the skull, age assessment of subadult remains is best done through a coordinated evaluation of such features as dentition and fontanelle closure, as well as radiographs and/or computed tomography (CT) scans. This is particularly key for studies of tooth development (calcification, eruption). It is important to emphasize that the evaluation of a skull without these methods is artificial and not reflective of actual practice. However, the ability to analyze such remains from the strict perspective of osteology is fundamental, and students must feel comfortable analyzing subadult skulls and skeletons.
4. It is not currently possible to reliably differentiate amongst the major racial groups within subadults.[7]
5. It is not currently possible to reliably differentiate male and female infant and young child skeletal remains.[7]
6. In the evaluation of subadult skulls, particularly when studying 'typical' eruption patterns, students must be cautioned that statistical data is based on **populations**, and may not necessarily be reflective of reality in an **individual**.
7. It may be appropriate to discuss the concept of sutural (Wormian) bones and what role they may play in the forensic evaluation of cranial remains. It is most important that students understand sutural bones are normal variants which may be present with somewhat increased frequency in some racial groups; they must not be misdiagnosed as fractures.

REFERENCES:

1. Kiesler, J. and Ricer, R. (2003). The abnormal fontanel. *American Family Physician*, 67(12): 2547-52.
2. Tunnessen, W.W., Jr. (1990). Persistent open anterior fontanelle. *JAMA*, 264(18): 2450.
3. Standring, S., Ed. (2005). *Gray's Anatomy: The Anatomical Basis of Clinical Practice*. 39 ed. London: Elsevier (Churchill Livingstone).
4. Konie, J. (1964). Comparative value of X-rays of the spheno-occipital synchondrosis and of the wrist for skeletal age assessment. *The Angle Orthodontist*, 34(4): 303-313.
5. Powell, T.V. and Brodie, A.G. (1963). Closure of the Spheno-Occipital Synchondrosis. *Anatomical Record*, 147: 15-23.
6. Scheuer, L. and Black, S. (2000). *Developmental Juvenile Osteology*. San Diego, CA: Elsevier (Academic Press).
7. Matshes, E. and Lew, E. (2006). Forensic osteology. In *Forensic Pathology: Principles and Practice*, D. Dolinak, E. Matshes, and E. Lew, Editors. San Diego, CA: Elsevier (Academic Press).

Bone Clones® Descriptive Analysis Report

DISCLAIMERS:

This report is meant only as a teaching tool for introductory level students of the anatomical, anthropology or forensic sciences who might be using this specimen to learn human and forensic osteology. Evaluation of osteologic material is best done with original specimens. My evaluation was based solely upon studies of a Bone Clones® replica. My opinions are based solely upon the material presented to me. This is somewhat artificial as in real forensic investigations additional studies would be undertaken prior to the formulation of diagnoses, and the production of a report. These studies might include plain film radiography, computed tomography (CT) studies, histology, etc. Evaluation of a child skull for age always involves radiography. Osteologic/odontologic evaluation of age based purely on visible eruption patterns is a useful basic skill, but is artificial and not representative of actual practice. My opinions regarding this skull were made without access to the postcranial skeleton.

Evan Matshes BSc, MD
Consultant Osteologist